



## OPEN ACCESS

## EDITED BY

Jonasz Jeremiasz Weber,  
Ruhr University Bochum, Germany

## REVIEWED BY

Argel Estrada-Mondragon,  
Linköping University, Sweden  
Tania López-Hernández,  
University of Barcelona, Spain

## \*CORRESPONDENCE

Sonja Sucic  
✉ sonja.sucic@meduniwien.ac.at

<sup>†</sup>These authors have contributed equally to this work

RECEIVED 18 July 2024

ACCEPTED 19 August 2024

PUBLISHED 29 August 2024

## CITATION

Shah N, Kasture AS, Fischer FP, Sitte HH, Hummel T and Sucic S (2024) A transporter's doom or destiny: *SLC6A1* in health and disease, novel molecular targets and emerging therapeutic prospects. *Front. Mol. Neurosci.* 17:1466694. doi: 10.3389/fnmol.2024.1466694

## COPYRIGHT

© 2024 Shah, Kasture, Fischer, Sitte, Hummel and Sucic. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

# A transporter's doom or destiny: *SLC6A1* in health and disease, novel molecular targets and emerging therapeutic prospects

Nikita Shah<sup>1†</sup>, Ameya S. Kasture<sup>1†</sup>, Florian P. Fischer<sup>2</sup>, Harald H. Sitte<sup>1,3,4</sup>, Thomas Hummel<sup>5</sup> and Sonja Sucic<sup>1\*</sup>

<sup>1</sup>Institute of Pharmacology, Medical University of Vienna, Vienna, Austria, <sup>2</sup>Department of Epileptology and Neurology, RWTH Aachen University, Aachen, Germany, <sup>3</sup>Hourani Center for Applied Scientific Research, Al-Ahliyya Amman University, Amman, Jordan, <sup>4</sup>Center for Addiction Research and Science-AddResS, Medical University of Vienna, Vienna, Austria, <sup>5</sup>Department of Neuroscience and Developmental Biology, University of Vienna, Vienna, Austria

As the first member of the solute carrier 6 (SLC6) protein family, the  $\gamma$ -aminobutyric acid (GABA) transporter 1 (GAT1, *SLC6A1*), plays a pivotal role in the uptake of GABA from the synaptic cleft into neurons and astrocytes. This process facilitates the subsequent storage of GABA in presynaptic vesicles. The human *SLC6A1* gene is highly susceptible to missense mutations, leading to severe clinical outcomes, such as epilepsy, in the afflicted patients. The molecular mechanisms of *SLC6A1*-associated disorders are discerned to some degree; many *SLC6A1* mutations are now known to impair protein folding, and consequently fail to reach the plasma membrane. Inherently, once inside the endoplasmic reticulum (ER), GAT1 abides by a complex cascade of events that enable efficient intracellular trafficking. This involves association with specialized molecular chaperones responsible for steering the protein folding process, oligomerization, sorting through the Golgi apparatus, and ultimately delivery to the cell surface. The entire process is subject to stringent quality control mechanisms at multiple checkpoints. While the majority of the existing loss-of-function *SLC6A1* variants interfere with folding and membrane targeting, certain mutants retain abundant surface expression. In either scenario, suppressed GAT1 activity disrupts GABAergic neurotransmission, preceding the disease manifestation in individuals harboring these mutations. The nervous system is enthralled and calls for systematic, groundbreaking research efforts to dissect the precise molecular factors associated with the onset of complex neurological disorders, and uncover additional non-canonical therapeutic targets. Recent research has given hope for some of the misfolded *SLC6A1* variants, which can be salvaged by small molecules, i.e., chemical and pharmacological chaperones, acting on multiple upstream targets in the secretory pathway. We here highlight the significance of pharmacochaperoning as a therapeutic strategy for the treatment of *SLC6A1*-related disorders.

## KEYWORDS

*SLC6A1*, GABA transporter 1, human disease variants, cellular quality control mechanisms, protein trafficking, epilepsy, intellectual disability, protein misfolding

## The ABC's of SLC6A1: basic physiology and structural features

The inhibitory neurotransmitter  $\gamma$ -aminobutyric acid (GABA) is essential for the regulation and function of various cortical and sub-cortical circuits. The right balance between excitation and inhibition in neural circuits is physiologically absolutely vital, and can lead to seizures when imbalanced (Treiman, 2001). GABA, released from the GABAergic neurons, is rapidly retrieved by the designated GABA transporters (GATs), expressed in GABAergic neurons and glia (Fattorini et al., 2020). Among the four GAT isoforms belonging to the solute carrier 6 (SLC6) family, GAT1 (SLC6A1) is the major GABA transporter in the central nervous system. It is responsible for clearing GABA from the extracellular space, thus establishing the basis for GABAergic signaling that terminates neurotransmission. The translocation of GABA via GAT1 depends on an inward electrochemical gradient, involving a co-transport of  $\text{Na}^+$  and  $\text{Cl}^-$  across the membrane (Scimemi, 2014). The GAT1 has 12 transmembrane domains in helical stretches, intracellular amino (N)- and carboxyl (C)-termini, and three glycosylation sites in its large second extracellular loop (Bennett and Kanner, 1997; Motiwala et al., 2022). Fluorescence resonance energy transfer (FRET) studies in HEK293 cells revealed that GAT1 exists as an assembly of multiple monomers (Schmid et al., 2001), with each

moiety capable of independent GABA uptake, as shown in *Xenopus laevis* oocytes (Soragna et al., 2005). Oligomerization also serves as a prerequisite for the Sec24D-dependent concentrative export of GAT1 from the endoplasmic reticulum (ER) compartment (Farhan et al., 2006).

## Cellular mechanisms of GAT1 synthesis and delivery to the plasmalemma

Being a transmembrane protein, GAT1 is largely hydrophobic, and its biogenesis machinery is similar to other secretory proteins, i.e., it comprises a series of steps dependent on the recognition of a hydrophobic peptide segment of the newly synthesized protein via the signal recognition particle (SRP). This recognition simultaneously slows down the translation process in the ribosomes (Korkhov et al., 2008). Subsequently, the hydrophobic peptide is directed towards the ER for insertion into the ER membrane via the Sec61 translocon complex, followed by the resumption of the translation process by ribosomes (Figure 1). As the translation of eukaryotic proteins is rather slow, protein folding takes place co-translationally (Balchin et al., 2016); i.e., protein synthesis and the insertion of peptide segments into the ER membrane occur concomitantly. During folding,




FIGURE 1

A detailed portrayal of GAT1 protein folding and membrane trafficking. Ribosome-translated nascent polypeptides are inserted into the ER membrane via the Sec61 translocon. The translocon provides a channel that shields the hydrophobic residues of the nascent chain, identifies transmembrane domains and hydrophobic signal peptides, and facilitates their lateral exit into the lipid bilayer. After lateral exit and N-linked glycosylation, transporters undergo calnexin/calreticulin cycles. Malectin, associated with ribophorin-I, also interacts with folding intermediates before the calnexin cycles begin, at sites distinct from the calnexin binding site in the ER and Golgi. HSP relay binds to the folding protein from the cytoplasmic side. Misfolded proteins or folding-intermediates, bound to chaperones, adhere to the ER-associated degradation (ERAD) pathway. Disengagement of HSP relay allows the engagement of COPII components with the natively folded oligomerized transporters, i.e., Sec24D interacts with the ER export motif located on the C-terminal of GAT1. From the ER-Golgi intermediate complex (ERGIC), GAT1 cargo is further transported to and through the Golgi complex. Exocyst and syntaxin 1A are recruited for proper targeting and insertion of GAT1 oligomers at the plasma membrane. Chemical and pharmacological chaperones can correct the folding of ER-retained/misfolded transporter mutants and rescue their expression (and activity) at the cell membrane.

the proteins undergo post-translational modifications, such as N-glycosylation and disulfide bond formation (Caramelo and Parodi, 2015). Molecular chaperone systems, like lectin-chaperones (e.g., calnexin, calreticulin, and malectin) and ATP-dependent molecular chaperones (e.g., heat shock proteins, HSPs) assist the folding process. This occurs in a highly coordinated manner, such that the glycoprotein undergoes numerous calnexin cycles, i.e., re-association of the protein with calnexin to achieve a natively folded structure, until it is no longer recognizable by calnexin, which in turn, prepares the protein cargo for ER exit (Booth and High, 2004).

The ER of eukaryotic cells is a very crowded place: the folding of large proteins can be inefficient and result in folding intermediates and misfolded proteins (Figure 1). Missense mutations, in, e.g., *SLC6* transporters, generally complicate this process even further. Proteins that fail to achieve a native state are removed from the system via ER-associated degradation (ERAD; Booth and High, 2004). The natively folded GAT1 forms homo-oligomers before exiting the ER (Scholze et al., 2002). Oligomeric GAT1 recruits Sec24D [a component of the coat protein complex II (COPII)] for concentrative ER export, which plays a key role in determining GAT1 levels at the plasma membrane (Farhan et al., 2007). Mutating the ER export motif on the C-terminus of GAT1, impedes its interaction with Sec24D, and exerts a dominant negative effect on wild-type GAT1 expression at the cell surface (Farhan et al., 2007). Similar effects were recently observed for a related SLC6 transporter, creatine transporter 1 (CRT1, *SLC6A8*), upon elimination of its N-terminal ER export motif (Ün et al., 2024).

COPII vesicles bud off the ER to form membrane-like fused structures termed the ER-Golgi intermediate compartment (ERGIC), from which GAT1 is further channeled toward the Golgi via a specific C-terminal motif (Farhan et al., 2008) for intra-Golgi trafficking (Figure 1). Additional post-translational modifications take place in the Golgi, which remove three residual mannose residues and adds complex molecules, like N-acetylglucosamines, fucose, galactose and sialic acid residues, which are required for mature glycosylation and priming the protein for the plasma membrane (Mohan et al., 2023). In essence, the proteins traverse through the cisternal membranes of the Golgi from *cis* to *trans* compartments, which are the key functional elements of the Golgi, with differential resident processing enzymes required for glycosylation (Jackson, 2009). The trafficking of secretory proteins from the *trans*-Golgi compartment to membrane surfaces remains poorly understood. Tubular pleomorphic structures budding from the *trans*-Golgi compartment are believed to be the carriers that transport secretory proteins directly to the plasma membrane (Stalder and Gershlick, 2020). The exocyst complex, however, is a well-known requirement for the regulated membrane insertion of the GAT1 (Farhan et al., 2004). The last three amino acids of GAT1 (AYI) interact with the exocyst (Farhan et al., 2004). The removal or substitution of these residues affects the membrane insertion of the transporters in cultured cells (Farhan et al., 2004).

Surface GAT1 levels are also highly orchestrated by other proteins, including the soluble N-ethylmaleimide-sensitive factor (NSF) attachment protein receptors (SNAREs), syntaxin-1A and synaptosomal-associated protein 25 (SNAP-25; Quick et al., 2004; Quick, 2006). Syntaxin-1A interacts with the N-terminus of GAT1 and increases its surface expression (Deken et al., 2000). Conversely, syntaxin-1A also acts as a negative regulator of GABA translocation, as the binding of syntaxin-1A to the N-terminus of GAT1 inhibits the flux, which can be reversed by GABA, due to its dissociation from syntaxin-1A (Quick, 2006). Thus, syntaxin-1A plays a dual role in tuning extracellular

neurotransmitter levels by regulating GAT1 surface expression via trafficking, and by modulating GABA transport. Syntaxin-1A is also involved in the docking and fusion of GABA-containing vesicles, suggesting that it regulates GABA release and reuptake. Interestingly, missense mutations in the gene encoding syntaxin-1A are also associated to epilepsy (Luppe et al., 2023).

## SLC6A1 loss-of-function pathogenic variants

Mutations in the *SLC6A1* gene, which encodes GAT1, have been associated with a wide range of neurodevelopmental disorders, such as autism, intellectual disability, and epilepsy syndromes (Fischer et al., 2022). The loss-of-function phenotype imparted by the mutations are attributed to single nucleotide variants or missense mutations, frameshift (fs)-, termination codon-and splice site-mutations (Figure 2). The *SLC6A1* gene is more sensitive to disruption by missense mutations than many other genes (Silva et al., 2024). Recurrent missense mutations (red boxes in Figure 2) are located at hypermutable CpG sites (Silva et al., 2024). These disease-linked missense mutations can be broadly classified into two groups, based on defective protein folding/trafficking or functional activity of the resulting GATs (Figure 2). Mutations that affect GABA transport, but not trafficking, typically impair the substrate binding site or its translocation pathway (blue spheres in Figure 2). In contrast, folding/trafficking-deficient variants are retained at the ER, but may exhibit substrate binding affinity similar to the wild-type protein (Figure 2, orange spheres), and are amenable to rescue by pharmacological means (green spheres in Figure 2).

## Digging deep in the ER: hunting for novel molecular and drug targets

SLC6 transporters undergo numerous conformational changes to reduce the free energy of folding and to attain a natively folded state. This is assisted by an elegant cascade of molecular chaperones and quality control mechanisms, described above. Point mutations in *SLC6A1* often disrupt the habitual folding trajectory of the transporter, trapping it in a misfolded state at the ER level. ER-resident chaperones play a crucial role in assisting such transporters (Korkhov et al., 2008; Kasture et al., 2016). The lectin-based luminal chaperone calnexin (Figure 1), can act as a “folding sensor” (Ou et al., 1993); e.g., it was shown to bind an oligomerization-deficient E101D-GAT1 mutant with markedly higher affinity than the wild-type (Korkhov et al., 2008). Similar observations were made for misfolded mutants in the transporters for dopamine (DAT, *SLC6A3*) and serotonin (SERT, *SLC6A4*; El-Kasaby et al., 2010; Koban et al., 2015; Kasture et al., 2016).

Malectin, a stress-responsive membrane-anchored-ER-resident lectin is responsible for the quality control of glycoproteins (Schallus et al., 2008, 2010; Qin et al., 2012; Feng et al., 2022). Malectin preferentially binds to misfolded proteins and inhibits their ER export (Qin et al., 2012). It does not compete with calnexin for substrate binding, suggesting an additional layer of lectin-based quality control for glycoproteins (Galli et al., 2011). Considering that malectin does not have an ER-retention peptide of its own, it



“trafficking-related players.” The GAT1 itself strides along the Golgi path. Since many epilepsy-linked GAT1 variants are stalled in the ER compartment (Kasture et al., 2023), it is feasible that the ensuing ER stress propagates, with an aftermath on Golgi dynamics. A recent study linked Golgi fragmentation with seizures in epileptic patients, as well as in *in vivo* studies in rats (Skupien-Jaroszek et al., 2023).

## The long and winding road from molecular mechanisms of disease to the rescue

The molecular chaperone system in cells facilitates proteins in achieving their thermodynamically stable and natively folded states. In the instance of misfolded GAT1 and other SLC6 transporters, this internal chaperone machinery falls short. However, folding-deficient mutants can be rescued by small molecules, i.e., chemical and pharmacological chaperones. Examples of compounds that have been effective in improving the expression and activity of such SLC6 variants, either *in vitro* or *in vivo*, are listed in Table 1. Chemical chaperones [e.g., glycerol, 4-PBA and dimethyl sulfoxide (DMSO)] are non-specific, and act on a wide range of proteins to promote their folding and delivery to the cell surface. Conversely, pharmacological chaperones (pharmacochaperones) display substrate specificity and bind and stabilize the target protein (Loo and Clarke, 2007). The folding trajectory of SLC6 transporters progresses through the inward-facing conformation of the transport cycle (Sucic et al., 2016; Kasture et al., 2017). Drugs that

bind and stabilize the inward-facing state of DAT and SERT can thus restore the folding/trafficking of these transporters (Kasture et al., 2016; Bhat et al., 2017; Mazhar Asjad et al., 2017). One such compound is noribogaine, which effectively restored the surface expression of misfolded transporters in cell lines and in living fruit flies (Kasture et al., 2016; Bhat et al., 2017; Mazhar Asjad et al., 2017). Tiagabine, a clinically approved antiseizure drug, locks GAT1 in its inward-open conformational state (Motiwala et al., 2022). In HEK293 cells expressing the GAT1 epilepsy-linked A288V variant (one of the most commonly recurrent mutations identified in patients; cf., Figure 2), we found a rather modest increase in uptake upon exposure to tiagabine. Surprisingly, in transgenic flies, the same treatment yielded significant increases in the surface expression of both wild-type GAT1 and A288V. Despite this, tiagabine treatment failed to reduce seizures in flies harboring the A288V mutation (Kasture et al., 2023). However, unsurpassed rescue was observed upon treatment with the chemical chaperone 4-PBA (Kasture et al., 2023). In addition to the role of 4-PBA in preventing the aggregation of misfolded proteins, it is also known to reduce ER stress (Kubota et al., 2006) and downregulate HSP70 activity (Rubenstein and Lyons, 2001). Since this chaperone modulates a wide range of activities at the cellular and molecular levels, it likely enhances the trafficking process in more ways than one. Previously, we showed that 4-PBA rescues folding-deficient CRT1 mutants, associated with the debilitating creatine transporter deficiency syndrome (El-Kasaby et al., 2019; Farr et al., 2020). SLC6 transporters are receptive to assimilating different substrates and co-substrates, allowing for flexibility, thus making them excellent drug targets (Bhat et al., 2021).

TABLE 1 Small molecules demonstrating successful restoration of cell surface expression and uptake activity of misfolded SLC6 transporter variants.

SLC6 member	Compounds with rescue potential	Misfolded variants	<i>In vitro/in vivo</i> study type	References
hDAT (SLC6A3)	Noribogaine and pifithrin- $\mu$	V158E, G327R, L368Q	Cell culture; humanized <i>Drosophila</i>	Mazhar Asjad et al. (2017)
	Pifithrin- $\mu$	L368Q	Patient-derived iPSCs	Ng et al. (2021)
	Bupropion	A314V, R445C	Cell culture	Beerepoot et al. (2016)
	Ibogaine analog 9b	A314V, G386R, P395L, R445C, R521W, P554L	Cell culture; humanized <i>Drosophila</i>	Bhat et al. (2021)
hSERT (SLC6A4)	Ibogaine and DMSO	RI(607,608)AA, RII(607-609)AAA	Cell culture	El-Kasaby et al. (2010)
	Noribogaine, pifithrin- $\mu$ , and 17-DMAG	PG(601,602)AA, F604Q, RI(607,608)AA	Cell culture	El-Kasaby et al. (2014)
	Phenethylamine (PAL)-287, -1,045, -1,046	PG(601,602)AA	Cell culture	Bhat et al. (2017)
	Ibogaine, noribogaine, ibogaine analogs 3b, 3c, 4c, 9a, 9b, 9c, 9d, PAL-287, -1,045, -1,046 and bupropion	PG(601,602)AA	Cell culture	Bhat et al. (2021)
	Trazodone, nefazodone, and vilazodone	PG(601,602)AA	Cell culture; humanized <i>Drosophila</i>	El-Kasaby et al. (2024)
hCRT1 (SLC6A8)	4-PBA	R391W, A404P, G424D, V539I, P544L, P554L	Cell culture	El-Kasaby et al. (2019)
hGlyT2 (SLC6A5)	4-PBA	Rescue of WT GlyT2 from the dominant negative S512R variant	Primary neurons	Arribas-González et al. (2015)
hGAT1 (SLC6A1)	4-PBA, liothyronine, and tiagabine	A288V, A334P, G362R, Y445C	Cell culture; humanized <i>Drosophila</i>	Kasture et al. (2023)

## Shaken but not stirred: acumens from recent studies in humanized flies

Targeted trafficking of transporter proteins is essential in both asymmetric cells like neurons, as well as in cells with complex morphologies, such as astrocytes. *In vivo* models allow us to study protein folding and trafficking in a physiological context. Due to their short life cycle, extensive genetic resources, ease of maintenance, and cost-effectiveness, *Drosophila melanogaster* has become a widely used animal model in studies of various aspects of neurobiology and neurological disorders (Kasture et al., 2018; Fischer et al., 2022, 2023). About 83% of all SLC superfamily of transporters are conserved between humans and flies (Ceder and Fredriksson, 2022). Most of the genes involved in the folding and quality control of proteins are also conserved across species. As a matter of fact, the HSP genes were first discovered in *Drosophila* (Ritossa, 1962). Mendelian or monogenic diseases are widely studied in flies. Various aspects of human proteins can be studied in a temporal and spatial manner by expressing them in fruit flies (Rubin and Spradling, 1982; Brand and Perrimon, 1993; Gratz et al., 2015; Mazhar Asjad et al., 2017; Kasture et al., 2019). We and others have studied disease-relevant human *SLC6* transporters and ion channels in fruit flies (Kasture et al., 2016, 2017, 2018, 2019; Sucic et al., 2016; Campbell et al., 2019; Hussain et al., 2024). *Drosophila* has a single GABA transporter (dGAT), expressed primarily in astrocytes (Stork et al., 2014). In addition to expressing dGAT, astrocytes also express a transporter for glutamate (Soustelle et al., 2002), thus regulating both extracellular GABA and glutamate levels. Knockout of the gene encoding dGAT results in embryonic lethality, while astrocytic knockdown induces locomotor defects in larvae and adult flies (Stork et al., 2014). A viable hypomorphic dGAT mutant, with a truncated N-terminus showed markedly reduced expression, increased sleep duration, but reduced sleep latency (Chaturvedi et al., 2022). Membrane expression of dGAT in astrocytes is highly dynamic and regulated by metabotropic GABA receptors (Muthukumar et al., 2014). Increased intracellular Ca<sup>2+</sup> concentrations in astrocytes lead to rapid endocytosis of dGAT, which results in suppression of neuronal activity (Zhang et al., 2017).

To understand the *in vivo* aspects of GAT1-linked disorders, we generated transgenic flies expressing YFP-tagged A288V-GAT1. These flies expressed both, human GAT1 carrying the A288V mutation, as well as the endogenous dGAT, and presented no recognizable phenotypic defects. The GAT1-A288V variant was trapped in the ER of transfected HEK293 cells, and was comparably contained within the ER compartment in A288V fly astrocytes (Kasture et al., 2023). When expressed in GABAergic neurons, A288V was restricted to the cell bodies (which contain the ER), while the wild-type GAT1 appeared to have a widespread expression pattern (Kasture et al., 2023). The dGAT was shown to modulate seizure behavior in flies (Muthukumar et al., 2014; Mazaud et al., 2019). Accordingly, we observed heat-induced seizures in A288V flies, recapitulating the clinical manifestations in people carrying the A288V mutation (Kasture et al., 2023). Interestingly, knockdown of GAT1 did not trigger heat-induced seizures in flies. In HEK293 cells, treatment with 4-PBA, liothyronine and tiagabine partly restored GABA uptake by A288V mutant. Our earlier studies on DAT taught us that even marginal rescue by pharmacological compounds *in vitro* can have profound physiological effects in *in vivo* systems (Kasture et al., 2016; Mazhar Asjad et al., 2017). In *Drosophila*, treatment with

4-PBA, liothyronine and tiagabine significantly improved the astrocytic expression of A288V in the antennal lobe neuropile of the adult fly brain, mimicking the wild-type GAT1 expression pattern and reduced seizure-like activity (Kasture et al., 2023), illuminating the pharmacochaperoning potential of these drugs, or congeners thereof, *in vivo*. Transgenic flies provide an efficient platform for exploring individual GAT1 pathogenic variants *in vivo*, and most importantly, testing the rescue capacity of various small molecules by, e.g., pharmacochaperoning.

## The existing treatment options for *SLC6A1*-related disorders

At present, there is no specific therapy available to target the underlying pathophysiologic mechanisms of *SLC6A1*-related disorders. Standard treatment is, hence, suboptimal and based on the presenting epilepsy syndrome, rather than the pertinent genetic etiology. Typically, broad spectrum antiseizure medications (ASMs) are used to manage the symptoms. A selection of the most widely used ASMs, in the context of *SLC6A1*-linked conditions, is listed in Table 2. In 2018, Johannesen and colleagues described a cohort of 34 patients harboring mutations in the *SLC6A1* gene, of which 31 had epilepsy (Johannesen et al., 2018). In this cohort, the most common seizure types were absence, myoclonic and atonic seizures. Around half of the patients were diagnosed with epilepsy accompanied with myoclonic-tonic seizures (EMAtS). Two thirds of the patients became seizure free under treatment with ASMs. Ten of these patients took valproic acid either as monotherapy or in combination with clonazepam, lamotrigine or levetiracetam. The remaining individuals became seizure free under treatment with ethosuximide, levetiracetam, lamotrigine or clonazepam, alone or in combination. Vagus nerve stimulation was attempted in one patient with Lennox–Gastaut syndrome without any beneficial effect. Importantly, the authors noted that there was no correlation between seizure control and cognitive outcome. In a recent case report, two further patients with isolated absence seizures were described (Caputo et al., 2023). Both patients became seizure free on valproic acid alone, or in combination with ethosuximide, respectively. Several further case reports described sound responses to valproic acid (Posar and Visconti, 2019; Poliquin et al., 2021; Mermer et al., 2022; Mori et al., 2023; Piniella et al., 2023). The favorable effect of valproic acid in *SLC6A1*-related disorders is not surprising, since it is known to modulate the GABAergic system by increasing the synthesis of GABA and by decreasing its degradation (Chateauvieux et al., 2010).

Beside ASMs, dietary approaches have also been reported as potentially effective treatment options for patients with *SLC6A1*-related disorders. For instance, one variant carrier with EMAtS showed an excellent response to a ketogenic diet. The authors described a virtual cessation of seizures after the diet had been started (Palmer et al., 2016). The ketogenic diet was also described as partially effective in another *SLC6A1* variant carrier in a previous cohort (Carvill et al., 2015). Notably, the ketogenic diet is generally considered a highly effective treatment option for patients with EMAtS (Oguni et al., 2002; Caraballo et al., 2006; Bergqvist, 2012). As most studies focus on the pediatric population, information on the long-term outcome of patients with *SLC6A1*-related disorders is relatively limited. Interestingly, a recent study described the phenotypic spectrum of 15 adult patients with *SLC6A1*-related disorders (Johannesen et al., 2023).

TABLE 2 A list of commonly used ASMs in the treatment of patients with *SLC6A1*-related disorders.

ASM	Mechanism(s) of action	References
Clobazam	Allosteric modulation of GABA <sub>A</sub> receptors	Carvill et al. (2015), Kassabian et al. (2023), Piniella et al. (2023), Trivisano et al. (2023)
Clonazepam	Allosteric modulation of GABA <sub>A</sub> receptors	Carvill et al. (2015), Johannesen et al. (2018), Trivisano et al. (2023)
Ethosuximide	Blockade of calcium channels (T-type)	Carvill et al. (2015), Johannesen et al. (2018), Caputo et al. (2023), Kassabian et al. (2023), Trivisano et al. (2023)
Lamotrigine	Blockade of voltage-gated sodium channels	Carvill et al. (2015), Johannesen et al. (2018), Cai et al. (2019), Kassabian et al. (2023), Mori et al. (2023), Trivisano et al. (2023)
Levetiracetam	Modulation of synaptic vesicle protein 2A (SV2A)	Carvill et al. (2015), Johannesen et al. (2018), Mermer et al. (2022), Kassabian et al. (2023), Trivisano et al. (2023)
Topiramate	Allosteric modulation of GABA <sub>A</sub> receptors, blockade of AMPA receptors, blockade of voltage-gated sodium channels, inhibition of carbonic anhydrase	Carvill et al. (2015), Johannesen et al. (2018), Kassabian et al. (2023)
Valproic acid	Blockade of voltage-gated sodium channels, blockade of calcium channels (T-type), activation of glutamic acid decarboxylase, inhibition of GABA transaminase	Carvill et al. (2015), Johannesen et al. (2018), Cai et al. (2019), Posar and Visconti (2019), Kahen et al. (2021), Mermer et al. (2022), Caputo et al. (2023), Trivisano et al. (2023), Kassabian et al. (2023), Mori et al. (2023), Piniella et al. (2023), Caraballo et al. (2024)

In this cohort, epilepsy was prevalent in 11/15 patients and the most common seizure types were absence, myoclonic, atonic and bilateral tonic-clonic seizures. Unfortunately, only four patients became seizure-free. These patients were administered either lamotrigine, valproic acid or a combination of the two drugs.

The high rate of refractory seizures further highlights the vast and unmet clinical need for the development of more efficacious treatments for intractable pediatric conditions, such as *SLC6A1*-triggered epilepsy. A current UK-based pilot clinical trial [NCT05437393; Children's Adaptive Deep brain stimulation for Epilepsy Trial (CADET)]—is investigating the benefits of a medical device surgically fitted onto a patient's skull and attached to electrodes which implement deep brain stimulation (DBS). Recently, DBS applied on a teenager suffering from life-threatening epilepsy (due to a mutation in the voltage-gated sodium channel beta subunit 1, *SCN1B*), ensued an astonishing 80% reduction in both seizure frequency and severity. DBS targeting the thalamic region specifically is known to be helpful in managing refractory seizures irresponsive to regular pharmacological interventions (Fisher, 2023). Since GAT1 shows abundant expression in the thalamus, and normal functioning of the thalamic GAT1 is essential for controlling synchronous spike-and-wave discharges (Cope et al., 2009), it is conceivable that DBS holds potential in handling seizures originating from GAT1 mutations. In light of our recent work (Kasture et al., 2023), it is worth pointing out that an ongoing clinical trial (NCT04937062) focuses on the use of glycerol phenylbutyrate (Ravicti) in patients with *SLC6A1*- and syntaxin-binding protein 1 (*STXBP1*)-associated developmental and epileptic encephalopathy.

## Summary

In the era of genome sequencing, the identification of monogenic disease-associated mutations is surging worldwide. *SLC6A1* epitomizes a gene especially prone to missense mutations, some of which happen to occur even more frequently due to hypermutable CpG sites (Silva et al., 2024). The current clinical management of *SLC6A1* disorders is far from impeccable. There has been a progressive collective effort, by our group and others, to seek new

and more effective pharmacological (and other) means in the treatment of *SLC6*-related pathologies. The pharmacochaperoning approach, in particular, proved exceptionally advantageous in rectifying protein folding and activity of dozens of pathogenic variants among *SLC6* family members (Beerepoot et al., 2016; Kasture et al., 2016, 2023; Bhat et al., 2017, 2021, 2023; Mazhar Asjad et al., 2017; El-Kasaby et al., 2019). 4-PBA, which is FDA-approved for the treatment of urea cycle disorders since the 1980s, has shown great merit in recouping disease variants of hCRT1 (El-Kasaby et al., 2019) and hGAT1 (Kasture et al., 2023). Thorough studies of the molecular machineries in the ER and other organelles, along with the adjoined endogenous quality control processes, ought to disclose prominent new drug targets.

## Author contributions

NS: Writing – original draft. AK: Writing – original draft. FF: Writing – original draft. HS: Funding acquisition, Project administration, Writing – review & editing. TH: Writing – review & editing. SS: Supervision, Conceptualization, Funding acquisition, Investigation, Project administration, Resources, Software, Validation, Visualization, Writing – original draft, Writing – review & editing.

## Funding

The author(s) declare that financial support was received for the research, authorship, and/or publication of this article. The authors acknowledge the financial support from the Austrian Science Fund (FWF), project P35589 to HS and P36574 to SS.

## Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated

organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

## References

- Arribas-González, E., De Juan-Sanz, J., Aragón, C., and López-Corcuera, B. (2015). Molecular basis of the dominant negative effect of a glycine transporter 2 mutation associated with hyperekplexia. *J. Biol. Chem.* 290, 2150–2165. doi: 10.1074/JBC.M114.587055
- Azoulay-Ginsburg, S., Di Salvio, M., Weitman, M., Afri, M., Ribeiro, S., Ebbinghaus, S., et al. (2021). Chemical chaperones targeted to the endoplasmic reticulum (ER) and lysosome prevented neurodegeneration in a C9orf72 repeat expansion *Drosophila* amyotrophic lateral sclerosis (ALS) model. *Pharmacol. Rep.* 73, 536–550. doi: 10.1007/S43440-021-00226-2
- Balchin, D., Hayer-Hartl, M., and Hartl, F. U. (2016). In vivo aspects of protein folding and quality control. *Science* 353:aac4354. doi: 10.1126/SCIENCE.AAC4354
- Beerepoot, P., Lam, V. M., and Salahpour, A. (2016). Pharmacological chaperones of the dopamine transporter rescue dopamine transporter deficiency syndrome mutations in heterologous cells. *J. Biol. Chem.* 291, 22053–22062. doi: 10.1074/JBC.M116.749119
- Bennett, E. R., and Kanner, B. I. (1997). The membrane topology of GAT-1, a (Na<sup>+</sup> + Cl<sup>-</sup>)-coupled gamma-aminobutyric acid transporter from rat brain. *J. Biol. Chem.* 272, 1203–1210. doi: 10.1074/JBC.272.2.1203
- Bergqvist, A. G. C. (2012). Myoclonic astatic epilepsy and the use of the ketogenic diet. *Epilepsy Res.* 100, 258–260. doi: 10.1016/j.EPLEPSYRES.2011.04.009
- Bhat, S., El-Kasaby, A., Freissmuth, M., and Sucic, S. (2021). Functional and biochemical consequences of disease variants in neurotransmitter transporters: a special emphasis on folding and trafficking deficits. *Pharmacol. Ther.* 222:107785. doi: 10.1016/j.PHARMTHERA.2020.107785
- Bhat, S., Hasenhuetl, P. S., Kasture, A., Ali, E. K., Baumann, M. H., Blough, B. E., et al. (2017). Conformational state interactions provide clues to the pharmacochaperone potential of serotonin transporter partial substrates. *J. Biol. Chem.* 292, 16773–16786. doi: 10.1074/JBC.M117.794081
- Booth, P. J., and High, S. (2004). Polytopic membrane protein folding and assembly *in vitro* and *in vivo*. *Mol. Membr. Biol.* 21, 163–170. doi: 10.1080/09687680410001697215
- Brand, A. H., and Perrimon, N. (1993). Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. *Development* 118, 401–415. doi: 10.1242/DEV.118.2.401
- Cai, K., Wang, J., Eissman, J., Wang, J., Nwosu, G., Shen, W., et al. (2019). A missense mutation in SLC6A1 associated with Lennox-Gastaut syndrome impairs GABA transporter 1 protein trafficking and function. *Exp. Neurol.* 320:112973. doi: 10.1016/j.expneurol.2019.112973
- Campbell, N. G., Shekar, A., Aguilar, J. I., Peng, D., Navratna, V., Yang, D., et al. (2019). Structural, functional, and behavioral insights of dopamine dysfunction revealed by a deletion in SLC6A3. *Proc. Natl. Acad. Sci. USA* 116, 3853–3862. doi: 10.1073/pnas.1816247116
- Caputo, D., Franceschetti, S., Castellotti, B., Freri, E., Zorzi, G., Saletti, V., et al. (2023). Case report: SLC6A1 mutations presenting with isolated absence seizures: description of 2 novel cases. *Front. Neurosci.* 17:1219244. doi: 10.3389/FNINS.2023.1219244
- Caraballo, R. H., Cersósimo, R. O., Sakr, D., Cresta, A., Escobal, N., and Fejerman, N. (2006). Ketogenic diet in patients with myoclonic-astatic epilepsy. *Epileptic Disord.* 8, 151–155. doi: 10.1684/j.1950-6945.2006.tb00174.x
- Caraballo, R. H., Chacón, S., Touzon, M. S., Loos, M., and Juanes, M. (2024). De novo absence status epilepticus associated with the SLC6A1 gene in a pediatric patient. *Epileptic Disord.* 26, 233–235. doi: 10.1002/EPD2.20175
- Caramelo, J. J., and Parodi, A. J. (2015). A sweet code for glycoprotein folding. *FEBS Lett.* 589, 3379–3387. doi: 10.1016/j.FEBSLET.2015.07.021
- Carvill, G. L., McMahon, J. M., Schneider, A., Zemel, M., Myers, C. T., Saykally, J., et al. (2015). Mutations in the GABA transporter SLC6A1 cause epilepsy with myoclonic-atonic seizures. *Am. J. Hum. Genet.* 96, 808–815. doi: 10.1016/j.AJHG.2015.02.016
- Ceder, M. M., and Fredriksson, R. (2022). A phylogenetic analysis between humans and *D. melanogaster*: A repertoire of solute carriers in humans and flies. *Gene* 809:146033. doi: 10.1016/j.GENE.2021.146033
- Chateauvieux, S., Morceau, F., Dicato, M., and Diederich, M. (2010). Molecular and therapeutic potential and toxicity of valproic acid. *J. Biomed. Biotechnol.* 2010, 1–18. doi: 10.1155/2010/479364
- Chaturvedi, R., Stork, T., Yuan, C., Freeman, M. R., and Emery, P. (2022). Astrocytic GABA transporter controls sleep by modulating GABAergic signaling in *Drosophila* circadian neurons. *Curr. Biol.* 32, 1895–1908.e5. doi: 10.1016/j.CUB.2022.02.066
- Chiba, P., Freissmuth, M., and Stockner, T. (2014). Defining the blanks – Pharmacochaperoning of SLC6 transporters and ABC transporters. *Pharmacol. Res.* 83, 63–73. doi: 10.1016/j.PHRS.2013.11.009
- Cope, D. W., Di Giovanni, G., Fyson, S. J., Orbán, G., Errington, A. C., Lrincz, M. L., et al. (2009). Enhanced tonic GABA A inhibition in typical absence epilepsy. *Nat. Med.* 15, 1392–1398. doi: 10.1038/nm.2058
- Deken, S. L., Beckman, M. L., Boos, L., and Quick, M. W. (2000). Transport rates of GABA transporters: regulation by the N-terminal domain and syntaxin 1A. *Nat. Neurosci.* 3, 998–1003. doi: 10.1038/79939
- El-Kasaby, A., Boytsov, D., Kasture, A., Krumpal, G., Hummel, T., Freissmuth, M., et al. (2024). Allosteric inhibition and Pharmacochaperoning of the serotonin transporter by the antidepressant drugs trazodone and Nefazodone. *Mol. Pharmacol.* 106, 56–70. doi: 10.1124/MOLPHARM.124.000881
- El-Kasaby, A., Just, H., Malle, E., Stolt-Bergner, P. C., Sitte, H. H., Freissmuth, M., et al. (2010). Mutations in the carboxyl-terminal SEC24 binding motif of the serotonin transporter impair folding of the transporter. *J. Biol. Chem.* 285, 39201–39210. doi: 10.1074/JBC.M110.118000
- El-Kasaby, A., Kasture, A., Koban, F., Hotka, M., Asjad, H. M. M., Kubista, H., et al. (2019). Rescue by 4-phenylbutyrate of several misfolded creatine transporter-1 variants linked to the creatine transporter deficiency syndrome. *Neuropharmacology* 161:107572. doi: 10.1016/j.NEUROPHARM.2019.03.015
- El-Kasaby, A., Koban, F., Sitte, H. H., Freissmuth, M., and Sucic, S. (2014). A cytosolic relay of heat shock proteins HSP70-1A and HSP90β monitors the folding trajectory of the serotonin transporter. *J. Biol. Chem.* 289, 28987–29000. doi: 10.1074/JBC.M114.595090
- Farhan, H., Freissmuth, M., and Sitte, H. H. (2006). Oligomerization of neurotransmitter transporters: a ticket from the endoplasmic reticulum to the plasma membrane. *Handb. Exp. Pharmacol.* 175, 233–249. doi: 10.1007/3-540-29784-7\_12
- Farhan, H., Korkhov, V. M., Paulitschke, V., Dorostkar, M. M., Scholze, P., Kudlacek, O., et al. (2004). Two discontinuous segments in the carboxyl terminus are required for membrane targeting of the rat gamma-aminobutyric acid transporter-1 (GAT1). *J. Biol. Chem.* 279, 28553–28563. doi: 10.1074/JBC.M307325200
- Farhan, H., Reiterer, V., Korkhov, V. M., Schmid, J. A., Freissmuth, M., and Sitte, H. H. (2007). Concentrative export from the endoplasmic reticulum of the gamma-aminobutyric acid transporter 1 requires binding to SEC24D. *J. Biol. Chem.* 282, 7679–7689. doi: 10.1074/JBC.M609720200
- Farhan, H., Reiterer, V., Kriz, A., Hauri, H. P., Pavelka, M., Sitte, H. H., et al. (2008). Signal-dependent export of GABA transporter 1 from the ER-Golgi intermediate compartment is specified by a C-terminal motif. *J. Cell Sci.* 121, 753–761. doi: 10.1242/JCS.017681
- Farr, C. V., El-Kasaby, A., Freissmuth, M., and Sucic, S. (2020). The Creatine transporter unfolded: A knotty premise in the cerebral Creatine deficiency syndrome. *Front. Synaptic Neurosci.* 12:588954. doi: 10.3389/FNSYN.2020.588954
- Fattorini, G., Melone, M., and Conti, F. (2020). A reappraisal of GAT-1 localization in neocortex. *Front. Cell. Neurosci.* 14:9. doi: 10.3389/FNCEL.2020.00009
- Feng, H., Qiu, T., Yin, C., Zhao, X., Xu, G., Qi, L., et al. (2022). The Rice Malectin regulates plant cell death and disease resistance by participating in glycoprotein quality control. *Int. J. Mol. Sci.* 23:5819. doi: 10.3390/IJMS23105819
- Fischer, F. P., Karge, R. A., Weber, Y. G., Koch, H., Wolking, S., and Voigt, A. (2023). *Drosophila melanogaster* as a versatile model organism to study genetic epilepsies: an overview. *Front. Mol. Neurosci.* 16:1116000. doi: 10.3389/FNMOL.2023.1116000
- Fischer, F. P., Kasture, A. S., Hummel, T., and Sucic, S. (2022). Molecular and clinical repercussions of GABA transporter 1 variants gone amiss: links to epilepsy and developmental spectrum disorders. *Front. Mol. Biosci.* 9:834498. doi: 10.3389/FMOLB.2022.834498
- Fisher, R. S. (2023). Deep brain stimulation of thalamus for epilepsy. *Neurobiol. Dis.* 179:106045. doi: 10.1016/j.NBD.2023.106045
- Galli, C., Bernasconi, R., Soldà, T., Calanca, V., and Molinari, M. (2011). Malectin participates in a backup glycoprotein quality control pathway in the mammalian ER. *PLoS One* 6:e16304. doi: 10.1371/JOURNAL.PONE.0016304
- Gratz, S. J., Rubinstein, C. D., Harrison, M. M., Wildonger, J., and O'Connor-Giles, K. M. (2015). CRISPR-Cas9 genome editing in *Drosophila*. *Curr. Protoc. Mol. Biol.* 111:31.2.1–31.2.20. doi: 10.1002/0471142727.MB3102S111
- Hussain, R., Lim, C. X., Shaikat, Z., Islam, A., Caseley, E. A., Lippiat, J. D., et al. (2024). *Drosophila* expressing mutant human KCNT1 transgenes make an effective tool

- for targeted drug screening in a whole animal model of KCNT1-epilepsy. *Sci. Rep.* 14:3357. doi: 10.1038/S41598-024-53588-X
- Jackson, C. L. (2009). Mechanisms of transport through the Golgi complex. *J. Cell Sci.* 122, 443–452. doi: 10.1242/JCS.032581
- Johannesen, K. M., Gardella, E., Linnankivi, T., Courage, C., de Saint Martin, A., Lehesjoki, A. E., et al. (2018). Defining the phenotypic spectrum of SLC6A1 mutations. *Epilepsia* 59, 389–402. doi: 10.1111/epi.13986
- Johannesen, K. M., Nielsen, J., Sabers, A., Isidor, B., Kattentidt-Mouravieva, A. A., Zieglgänsberger, D., et al. (2023). The phenotypic presentation of adult individuals with SLC6A1-related neurodevelopmental disorders. *Front. Neurosci.* 17:1216653. doi: 10.3389/FNINS.2023.1216653
- Kahen, A., Kavus, H., Geltzeiler, A., Kentros, C., Taylor, C., Brooks, E., et al. (2021). Neurodevelopmental phenotypes associated with pathogenic variants in SLC6A1. *J. Med. Genet.* 59, 536–543. doi: 10.1136/jmedgenet-2021-107694
- Kassabian, B., Fenger, C. D., Willems, M., Aledo-Serrano, A., Linnankivi, T., McDonnell, P. P., et al. (2023). Intrafamilial variability in SLC6A1-related neurodevelopmental disorders. *Front. Neurosci.* 17:1219262. doi: 10.3389/FNINS.2023.1219262
- Kasture, A. S., Bartel, D., Steinkellner, T., Susic, S., Hummel, T., and Freissmuth, M. (2019). Distinct contribution of axonal and somatodendritic serotonin transporters in *Drosophila* olfaction. *Neuropharmacology* 161:107564. doi: 10.1016/j.neuropharm.2019.03.007
- Kasture, A., El-Kasaby, A., Szölösi, D., Mazhar Asjad, H. M., Grimm, A., Stockner, T., et al. (2016). Functional rescue of a misfolded *Drosophila melanogaster* dopamine transporter mutant associated with a sleepless phenotype by pharmacological chaperones. *J. Biol. Chem.* 291, 20876–20890. doi: 10.1074/JBC.M116.737551
- Kasture, A. S., Fischer, F. P., Kunert, L., Burger, M. L., Burgstaller, A. C., El-Kasaby, A., et al. (2023). *Drosophila melanogaster* as a model for unraveling unique molecular features of epilepsy elicited by human GABA transporter 1 variants. *Front. Neurosci.* 16:1074427. doi: 10.3389/FNINS.2022.1074427
- Kasture, A. S., Hummel, T., Susic, S., and Freissmuth, M. (2018). Big lessons from tiny flies: *Drosophila melanogaster* as a model to explore dysfunction of dopaminergic and serotonergic neurotransmitter systems. *Int. J. Mol. Sci.* 19:1788. doi: 10.3390/ijms19061788
- Kasture, A., Stockner, T., Freissmuth, M., and Susic, S. (2017). An unfolding story: small molecules remedy misfolded monoamine transporters. *Int. J. Biochem. Cell Biol.* 92, 1–5. doi: 10.1016/J.BIOCELL.2017.09.004
- Koban, F., El-Kasaby, A., Häusler, C., Stockner, T., Simbrunner, B. M., Sitte, H. H., et al. (2015). A salt bridge linking the first intracellular loop with the C terminus facilitates the folding of the serotonin transporter. *J. Biol. Chem.* 290, 13263–13278. doi: 10.1074/JBC.M115.641357
- Korkhov, V. M., Milan-Lobo, L., Zuber, B., Farhan, H., Schmid, J. A., Freissmuth, M., et al. (2008). Peptide-based interactions with calnexin target misassembled membrane proteins into endoplasmic reticulum-derived multilamellar bodies. *J. Mol. Biol.* 378, 337–352. doi: 10.1016/J.JMB.2008.02.056
- Kubota, K., Niinuma, Y., Kaneko, M., Okuma, Y., Sugai, M., Omura, T., et al. (2006). Suppressive effects of 4-phenylbutyrate on the aggregation of Pael receptors and endoplasmic reticulum stress. *J. Neurochem.* 97, 1259–1268. doi: 10.1111/J.1471-4159.2006.03782.X
- Loo, T. W., and Clarke, D. M. (2007). Chemical and pharmacological chaperones as new therapeutic agents. *Expert Rev. Mol. Med.* 9, 1–18. doi: 10.1017/S1462399407000361
- Luppe, J., Sticht, H., Lecoquierre, F., Goldenberg, A., Gorman, K. M., Molloy, B., et al. (2023). Heterozygous and homozygous variants in STX1A cause a neurodevelopmental disorder with or without epilepsy. *Eur. J. Hum. Genet.* 31, 345–352. doi: 10.1038/S41431-022-01269-6
- Mazaud, D., Kottler, B., Gonçalves-Pimentel, C., Proelss, S., Tüchler, N., Deneuborg, C., et al. (2019). Transcriptional regulation of the glutamate/GABA/glutamine cycle in adult glia controls motor activity and seizures in *Drosophila*. *J. Neurosci.* 39, 5269–5283. doi: 10.1523/JNEUROSCI.1833-18.2019
- Mazhar Asjad, H. M., Kasture, A., El-Kasaby, A., Sackel, M., Hummel, T., Freissmuth, M., et al. (2017). Pharmacochaperoning in a *Drosophila* model system rescues human dopamine transporter variants associated with infantile/juvenile parkinsonism. *J. Biol. Chem.* 292, 19250–19265. doi: 10.1074/JBC.M117.797092
- Mermer, F., Poliquin, S., Zhou, S., Wang, X., Ding, Y., Yin, F., et al. (2022). Astrocytic GABA transporter 1 deficit in novel SLC6A1 variants mediated epilepsy: connected from protein destabilization to seizures in mice and humans. *Neurobiol. Dis.* 172:105810. doi: 10.1016/J.NBD.2022.105810
- Mohan, A. G., Calenic, B., Ghiurau, N. A., Duncea-Borca, R. M., Constantinescu, A. E., and Constantinescu, I. (2023). The Golgi apparatus: A voyage through time, structure, function and implication in neurodegenerative disorders. *Cells* 12:1972. doi: 10.3390/CELLS12151972
- Mori, T., Sakamoto, M., Tayama, T., Goji, A., Toda, Y., Fujita, A., et al. (2023). A case of epilepsy with myoclonic atonic seizures caused by SLC6A1 gene mutation due to balanced chromosomal translocation. *Brain Dev.* 45, 395–400. doi: 10.1016/J.BRAINDEV.2023.03.001
- Motiwala, Z., Aduri, N. G., Shaye, H., Han, G. W., Lam, J. H., Katritch, V., et al. (2022). Structural basis of GABA reuptake inhibition. *Nature* 606, 820–826. doi: 10.1038/s41586-022-04814-x
- Muthukumar, A. K., Stork, T., and Freeman, M. R. (2014). Activity-dependent regulation of astrocyte GAT levels during synaptogenesis. *Nat. Neurosci.* 17, 1340–1350. doi: 10.1038/nn.3791
- Nakagomi, S., Barsoum, M. J., Bossy-Wetzell, E., Sütterlin, C., Malhotra, V., and Lipton, S. A. (2008). A Golgi fragmentation pathway in neurodegeneration. *Neurobiol. Dis.* 29, 221–231. doi: 10.1016/J.NBD.2007.08.015
- Ng, J., Barral, S., de la Fuente Barrigon, C., Lignani, G., Erdem, F. A., Wallings, R., et al. (2021). Gene therapy restores dopamine transporter expression and ameliorates pathology in iPSC and mouse models of infantile parkinsonism. *Sci. Transl. Med.* 13:eaaw1564. doi: 10.1126/scitranslmed.aaw1564
- Oguni, H., Tanaka, T., Hayashi, K., Funatsuka, M., Sakauchi, M., Shirakawa, S., et al. (2002). Treatment and long-term prognosis of myoclonic-astatic epilepsy of early childhood. *Neuropediatrics* 33, 122–132. doi: 10.1055/S-2002-33675
- Ou, W. J., Cameron, P. H., Thomas, D. Y., and Bergeron, J. J. M. (1993). Association of folding intermediates of glycoproteins with calnexin during protein maturation. *Nature* 364, 771–776. doi: 10.1038/364771A0
- Palmer, S., Towne, M. C., Pearl, P. L., Pelletier, R. C., Genetti, C. A., Shi, J., et al. (2016). SLC6A1 mutation and ketogenic diet in epilepsy with myoclonic-atonic seizures. *Pediatr. Neurol.* 64, 77–79. doi: 10.1016/j.pediatrneuro.2016.07.012
- Piniella, D., Canseco, A., Vidal, S., Xiol, C., Díaz de Bustamante, A., Martí-Carrera, I., et al. (2023). Experimental and bioinformatic insights into the effects of epileptogenic variants on the function and trafficking of the GABA transporter GAT-1. *Int. J. Mol. Sci.* 24:955. doi: 10.3390/IJMS24020955
- Poliquin, S., Hughes, I., Shen, W., Mermer, F., Wang, J., Mack, T., et al. (2021). Genetic mosaicism, intrafamilial phenotypic heterogeneity, and molecular defects of a novel missense SLC6A1 mutation associated with epilepsy and ADHD. *Exp. Neurol.* 342:113723. doi: 10.1016/j.expneurol.2021.113723
- Posar, A., and Visconti, P. (2019). Mild phenotype associated with SLC6A1 gene mutation: a case report with literature review. *J. Pediatr. Neurosci.* 14, 100–102. doi: 10.4103/jpn.JPN\_2\_19
- Qin, S. Y., Hu, D., Matsumoto, K., Takeda, K., Matsumoto, N., Yamaguchi, Y., et al. (2012). Malectin forms a complex with ribophorin I for enhanced association with misfolded glycoproteins. *J. Biol. Chem.* 287, 38080–38089. doi: 10.1074/JBC.M112.394288
- Quick, M. W. (2006). The role of SNARE proteins in trafficking and function of neurotransmitter transporters. *Handb. Exp. Pharmacol.* 175, 181–196. doi: 10.1007/3-540-29784-7\_9
- Quick, M. W., Hu, J., Wang, D., and Zhang, H. Y. (2004). Regulation of a gamma-aminobutyric acid transporter by reciprocal tyrosine and serine phosphorylation. *J. Biol. Chem.* 279, 15961–15967. doi: 10.1074/JBC.M306924200
- Rajeshwari, B. R., Shah, N., Joshi, P., Madhusudan, M. S., and Balasubramanian, N. (2023). Kinetics of Arf1 inactivation regulates Golgi organisation and function in non-adherent fibroblasts. *Biol. Open* 12:bio059669. doi: 10.1242/BIO.059669
- Ritossa, F. (1962). A new puffing pattern induced by temperature shock and DNP in *Drosophila*. *Experientia* 18, 571–573. doi: 10.1007/BF02172188
- Rubenstein, R. C., and Lyons, B. M. (2001). Sodium 4-phenylbutyrate downregulates HSC70 expression by facilitating mRNA degradation. *Am. J. Phys. Lung Cell. Mol. Phys.* 281, L43–L51. doi: 10.1152/AJPLUNG.2001.281.1.L43
- Rubin, G. M., and Spradling, A. C. (1982). Genetic transformation of *Drosophila* with transposable element vectors. *Science* 218, 348–353. doi: 10.1126/SCIENCE.6289436
- Schallus, T., Fehér, K., Sternberg, U., Rybin, V., and Muhle-Goll, C. (2010). Analysis of the specific interactions between the lectin domain of malectin and diglycosides. *Glycobiology* 20, 1010–1020. doi: 10.1093/GLYCOB/CWQ059
- Schallus, T., Jaechk, C., Fehér, K., Palma, A. S., Liu, Y., Simpson, J. C., et al. (2008). Malectin: a novel carbohydrate-binding protein of the endoplasmic reticulum and a candidate player in the early steps of protein N-glycosylation. *Mol. Biol. Cell* 19, 3404–3414. doi: 10.1091/MBE08-04-0354
- Schmid, J. A., Scholze, P., Kudlacek, O., Freissmuth, M., Singer, E. A., and Sitte, H. H. (2001). Oligomerization of the human serotonin transporter and of the rat GABA transporter 1 visualized by fluorescence resonance energy transfer microscopy in living cells. *J. Biol. Chem.* 276, 3805–3810. doi: 10.1074/jbc.M007357200
- Scholze, P., Freissmuth, M., and Sitte, H. H. (2002). Mutations within an intramembrane leucine heptad repeat disrupt oligomer formation of the rat GABA transporter 1. *J. Biol. Chem.* 277, 43682–43690. doi: 10.1074/jbc.M205602200
- Scimemi, A. (2014). Structure, function, and plasticity of GABA transporters. *Front. Cell. Neurosci.* 8:161. doi: 10.3389/FNCEL.2014.00161
- Shi, Y., Wang, S., Wu, J., Jin, X., and You, J. (2021). Pharmaceutical strategies for endoplasmic reticulum-targeting and their prospects of application. *J. Control. Release* 329, 337–352. doi: 10.1016/J.JCONREL.2020.11.054
- Silva, D. B., Trinidad, M., Ljungdahl, A., Revalde, J. L., Berguig, G. Y., Wallace, W., et al. (2024). Haploinsufficiency underlies the neurodevelopmental consequences of SLC6A1 variants. *Am. J. Hum. Genet.* 111, 1222–1238. doi: 10.1016/J.AJHG.2024.04.021

- Singh, V., Erady, C., and Balasubramanian, N. (2018). Cell-matrix adhesion controls Golgi organization and function through Arf1 activation in anchorage-dependent cells. *J. Cell Sci.* 131:jcs215855. doi: 10.1242/JCS.215855
- Skupien-Jaroszek, A., Szczepankiewicz, A. A., Rysz, A., Marchel, A., Matyja, E., Grajkowska, W., et al. (2023). Morphological alterations of the neuronal Golgi apparatus upon seizures. *Neuropathol. Appl. Neurobiol.* 49:e12940. doi: 10.1111/NAN.12940
- Soragna, A., Bossi, E., Giovannardi, S., Pisani, R., and Peres, A. (2005). Functionally independent subunits in the oligomeric structure of the GABA cotransporter rGAT1. *Cell. Mol. Life Sci.* 62, 2877–2885. doi: 10.1007/s00018-005-5322-x
- Soustelle, L., Besson, M. T., Rival, T., and Birman, S. (2002). Terminal glial differentiation involves regulated expression of the excitatory amino acid transporters in the *Drosophila* embryonic CNS. *Dev. Biol.* 248, 294–306. doi: 10.1006/dbio.2002.0742
- Stalder, D., and Gershlick, D. C. (2020). Direct trafficking pathways from the Golgi apparatus to the plasma membrane. *Semin. Cell Dev. Biol.* 107, 112–125. doi: 10.1016/j.semcdb.2020.04.001
- Stork, T., Sheehan, A., Tasdemir-Yilmaz, O. E., and Freeman, M. R. (2014). Neuron-glia interactions through the heartless FGF receptor signaling pathway mediate morphogenesis of *Drosophila* astrocytes. *Neuron* 83, 388–403. doi: 10.1016/j.neuron.2014.06.026
- Sucic, S., Kasture, A., Asjad, H. M. M., Kern, C., El-Kasaby, A., and Freissmuth, M. (2016). When transporters fail to be transported: how to rescue folding-deficient SLC6 transporters. *J. Neurol. Neuromed.* 1, 34–40. doi: 10.29245/2572.942X/2016/9.1098
- Treiman, D. M. (2001). GABAergic mechanisms in epilepsy. *Epilepsia* 42, 8–12. doi: 10.1046/J.1528-1157.2001.042SUPPL.3008.X
- Trivisano, M., Butera, A., Quintavalle, C., De Dominicis, A., Calabrese, C., Cappelletti, S., et al. (2023). Insights into cognitive and behavioral comorbidities of SLC6A1-related epilepsy: five new cases and literature review. *Front. Neurosci.* 17:1215684. doi: 10.3389/FNINS.2023.1215684
- Ün, D., Kovalchuk, V., El-Kasaby, A., Kasture, A., Koban, F., Kudlacek, O., et al. (2024). Breaking the rules of SLC6 transporters: export of the human creatine transporter-1 from the endoplasmic reticulum is supported by its N-terminus. *J. Neurochem.* doi: 10.1111/JNC.16088 [Epub ahead of print].
- Yang, Q. P., Fu, M. F., Gao, H., Yamamoto, K., Hu, D., and Qin, S. Y. (2018). Subcellular distribution of endogenous malectin under rest and stress conditions is regulated by ribophorin I. *Glycobiology* 28, 374–381. doi: 10.1093/GLYCOB/CWY034
- Zhang, Y. V., Ormerod, K. G., and Littleton, J. T. (2017). Astrocyte  $Ca_2^+$  influx negatively regulates neuronal activity. *eNeuro* 4, ENEURO.0340–ENEU16.2017. doi: 10.1523/ENEURO.0340-16.2017

## Glossary

17-DMAG	17-Dimethylaminoethylamino-17-demethoxygeldanamycin
4-PBA	4-Phenylbutyric acid
ALS	Amyotrophic lateral sclerosis
Arf1	ADP-ribosylation factor1
ASM	Antiseizure medication
COPII	Coat protein complex II
CRT1	Creatine transporter-1
DAT	Dopamine transporter
DMSO	Dimethyl sulfoxide
EMAs	Epilepsy with myoclonic-atonic seizures
ER	Endoplasmic reticulum
ERAD	ER-associated degradation
ERGIC	ER-Golgi intermediate compartment
FRET	Fluorescence resonance energy transfer
GABA	$\gamma$ Aminobutyric acid
GAT1	GABA transporter-1
GTPases	Guanosine triphosphatases
HSP	Heat shock protein
iPSC	Induced pluripotent stem cells
KDEL	Lys-Asp-Glu-Leu
NSF	N-ethylmaleimide-sensitive factor
QC	Quality control
SERT	Serotonin transporter
SLC6	Solute carrier 6
SNAP-25	Synaptosomal-associated protein 25
SNARE	Soluble N-ethylmaleimide-sensitive factor attachment protein receptor
SRP	Signal recognition particle
SV2A	Synaptic vesicle protein 2A